

International Journal of Engineering, Agriculture and Medicine (IJEAM)

An International Peer Review E-3 Journal of Science Innovation Technology

Section A – Engineering Sciences

Section B – Agriculture Sciences

Section C – Medical Sciences

Available online at www.ijeam.webs.com

Research Article

ETHYLMETHANE SULPHONATE INDUCED BUSHY HEAD MUTANTS IN JUTE (*Corchorus olitorius* L. Variety JRO-632).

P.K.Ghosh^{*1} and A.Chatterjee²

¹CSB, CSR&TI, Berhampore-742101,
Murshidabad, West Bengal.
paritush.ghosh@rediffmail.com

²Centre of Advanced Study in Cell and Chromosome Research,
Department of Botany, University of Calcutta,
Calcutta-700047. West Bengal.
chatterjeeja@email.com

Abstract: Presoaked seeds of jute (*Corchorus olitorius* L. Variety JRO-632) were treated with 0.5% Ethylmethane Sulphonate (EMS) for 24 hours. Bushy head mutants were screened in M3 in contrast to the normal plants. Bushy head mutants otherwise looked normal excepting the nature of early flowering habit. A number of yield component growth parameters were recorded like plant height, basal diameter, plant spread, root length, pod per plant, seeds per pod, pod length/ breadth ratio, number of primary branches per plant, number of secondary branches per plant, leaf angle, branching angle, first flowering date, 100% flowering date, total duration, % of pollen sterility and weight of 100 seeds which were found to vary from the control plant. Chromosome analysis revealed a number of aberrations like stickiness, fragmentation, clumping, polyploidy, and laggard and bridge formation etc. at very low frequency. These early flowering mutant plants gives more fiber yield than the control plants with superior quality. Multiple cropping has been possible with the availability of irrigation water and a number of early maturing varieties have introduced in case of various other crops. There should be a suitable early maturing variety of jute also to be best fitted in the multiple cropping patterns. With this objective in view the work on induction of mutation with chemical mutagen Ethylmethane sulphonate (EMS) was initiated.

Keywords: Ethylmethane Sulphonate (EMS, *Corchorus olitorius* L., Bushy head Mutant, 24 hours, chromosome, concentration 0.5%.

1. INTRODUCTION

Jute (*Corchorus olitorius* L. Variety JRO-632) is one of the very important fiber yielding cash crops with great demand in International market. A number of mutants in jute were reported through genetic manipulation by application of ionizing radiations (Kundu, 1944, Ghosh, 1969, Hossain, 1970 and Basu, 1967, Bandhyopadhyay, 1974 and Monti, 1968). However. Chemical mutagenesis in jute is still lacking although considerable worked has been done on this line in other commercial crops, the present work was therefore, undertaken to investigate the potentiality of host chemical substances to induce mutation in jute.

2. MATERIALS AND METHODS

Jute seeds (*Corchorus olitorius* L. Variety JRO-632) obtained from Jute Agriculture Research Institute, ICAR, Barrackpore, W.B. were presoaked in distilled water for 24 hours and then treated with 0.5% Ethylmethane sulphonate for 24 hours. The seeds were thoroughly washed with distilled water and then sown in the field directly with equal spacing for raising M1 generation. The individual M1 plants were harvested separately for growing M2 generation in progeny rows. The bushy head mutants were screened and again harvested for raising M3 generation. The mutants were screened in M3 generation. A number of

essential yield components were recorded. Cytological anomalies and pollen sterility were recorded as per schedule techniques.

3. RESULTS AND DISCUSSION

The present investigation indicates that some bushy head mutants were screened after M3 generations. These bushy head mutants otherwise looked normal plants like. Variation of yield component growth parameters was recorded. The segregation behavior in M2 generations was fitted to a ratio 3:1. In M2 almost all the plants were 8-locular fruit mutants excepting one or two cases of normal plants. No much noticeable variation in chromosome anomalies was recorded. Segregation behavior indicates that this is due to a single gene of recessive nature and pollen sterility was also recorded. Multiple cropping has been possible with the availability of irrigation water and a number of bushy head varieties have introduced in case of various other crops. There should be a suitable early maturing variety of jute also to be best fitted in the multiple cropping patterns. With this objective in view the work on induction of mutation with chemical mutagen Ethylmethane sulphonate (EMS) was initiated.

References

- [1] Bandyopadhyay, B. (1974). Studies on induced mutation in jute (*Corchorus capsularies* L. and *C. olitorius* L.) by single and combined treatments with X-rays and chemicals. Ph.D. Thesis, University of Calcutta, Calcutta.
- [2] Basu, R. K. (1967). Induction of high hosh, N. (1969). Studies of variability yielding mutants in jute. *Mutat. Res.* 4:163-167.
- [3] Gin jute followed by irradiation and hybridization. Ph.D. Thesis, BCKV.
- [4] Hussein, M.M. (1970). Cytogenetic studies of mutants and colchipooids of some cultivated varieties of jute. Ph.D. Thesis, BCKV.
- [5] Kundu, B.C. (1944). A note on the preliminary studies on effect of X-irradiation upon growth and development of jute plant. *Sci. & Cult.*, 9:559-560.
- [6] Monti, L.M. (1968). Mutations in peas induced by Diethyl Sulphate and X-rays. *Mutat. Res.*, 5:187-191.